

## 8

---

## The Role of Context in Behavioral Effects of Foods

*Herbert L. Meiselman and  
F. Matthew Kramer<sup>1</sup>*

### INTRODUCTION

The situation in which we eat, the eating context, is receiving increasing attention as a major factor in the control of eating habits. Traditionally, more attention was placed on the food itself and the personality characteristics and physiology of the eater. Placing food in its context has led to research and discussion of a number of situational or contextual variables such as the effort needed to obtain food (Collier, 1989; Meiselman et al., 1988), the time of day food is eaten (Birch et al., 1984; Kramer et al., 1992), the presence of other food, and social dynamics (de Castro and Brewer, 1991). Schutz (1988) sought to capture the overall impact of eating contexts in his measurement of "appropriateness," which indicates how well a food fits into a particular situation.

The emerging role of context has been reflected in its inclusion in recent international meetings. At the First International Conference on Food Choice,

---

<sup>1</sup> Herbert L. Meiselman and F. Matthew Kramer, U.S. Army Natick, ATTN: SATNC-Y, Natick, MA 01760-5020

a multidisciplinary meeting held in Brussels, Belgium, in July 1992, a number of sessions were devoted to the cultural and social context of eating, and papers throughout the meeting addressed context as well. At the meeting on Advances in Sensory Science, held in Helsinki, Finland, in August 1992 in memory of Professor Rose Marie Pangborn, an entire session was devoted to context.

The U.S. Army's interest in context began with large field studies of the military ration, the Meal Ready to Eat, in the early 1980s. The traditional approach was to research and evaluate food product development and improvement. In the early 1980s, however, researchers began to realize that without attention to the eating situation they could never fully understand what controls eating and what should be done to maximize soldiers' intakes of needed nutrients. Some of that research is presented in articles by Meiselman et al. (1988) and Hirsch and Kramer (1993).

Not only was it necessary to rethink what controls eating, but it was also necessary to rethink how to conduct research. Researchers began to doubt the adequacy of short-term laboratory tests, and so they focused more on longer-term field tests that better covered the situational variables that are now viewed as critical. The questioning of how human food research is conducted led to a recent critical review of methodology in the journal *Appetite* (Meiselman, 1992).

The role of context has long been realized in the experimental design literature (Campbell and Stanley, 1966). The difference now may be that while context was traditionally seen as a source of confounding to be controlled, more effort is currently being put into defining and studying the impact of context. The change in research is not only in how research is conducted but also in the focus.

A useful example might be derived from the growing body of research on noncaloric substances such as sugar and fat substitutes as human foods. The initial studies were simple, looking at the impact of a sweet drink (e.g., aspartame-containing beverages) on the subsequent intake of a test meal. In principle, to the subjects the drinks or other vehicles were made to seem equivalent, so that any differences in response to the drinks could be attributed to the fact that one contained calories and the other did not. Among these early studies of aspartame were reports (Blundell and Hill, 1986) suggesting that consumption of such items enhanced hunger and possibly intake as well, implying that diet drinks would not only fail to produce useful reductions in caloric intake but might actually be associated with increases. Early studies also indicated that typical two-group comparisons could not readily separate out the role of calories versus the role of sweetness (Blundell et al., 1988). Given the nature of these findings, aspartame and other noncaloric sweeteners have been the subject of many recent studies (see Rolls [1991] for a review), but those studies have not yet clarified their impacts on food habits. What

those studies  
context of the  
with an increa  
(Rolls, 1991)  
administrative  
chewing gum  
measurement  
outcome mea  
ratings, there  
1 to 2 weeks  
that they we

The role of  
mood and the  
the context of  
the food and  
situation in v  
discuss the r  
performance-  
covered beca  
the reader of  
the effects of  
looking at fo  
researchers to

Studies of  
baseline cond  
Rapaport (19  
and help to co  
tests. However  
treatment ord

Christensen  
immediate pr  
consumed the  
from eating f  
before the test  
about what el  
is being chang  
patterns and c

those studies have made clear is that the results and conclusions reflect the context of the study. Depending on the investigation, aspartame is associated with an increase, decrease, or no change in eating or other outcome measures (Rolls, 1991). The context of the studies has varied across the means of administration (flavored or unflavored beverages, food and meal ingredients, chewing gum, capsule form), timing of administration (preload before a measurement period, during a meal or meals, at multiple times during the day), outcome measures (consumption, palatability, hunger or other subjective ratings, thermic effect of ingestion), length of study (single meals, 24-h impact, 1 to 2 weeks), and participant expectations (whether subjects were informed that they were taking aspartame, use of aspartame as a diet aid).

The role of context in studying the effects of foods on performance and mood and then applying those effects covers a wide area. One could discuss the context of the research situation, its physical and social environment, and the food and its immediate context. One could discuss the nonresearch eating situation in which, ultimately, foods of interest are consumed. One could also discuss the military context, which investigators are trying to address using performance- and mood-altering foods. Not all aspects of the topic can be covered because of the current state of the art, but this chapter should remind the reader of context and perhaps raise the reader's assessment of its role in the effects of food on performance and mood. Context can be covered by looking at foods, the people who eat them, and the environment in which researchers test and try to predict peoples responses.

#### DETERMINING A BASELINE

Studies of the effects of foods on performance or mood generally need a baseline condition against which treatment effects can be measured. Kruesi and Rapaport (1986) point out that baseline data can be used to estimate variance and help to control problems of individual differences, a key problem in these tests. However, Kruesi and Rapaport also note the potential problems of treatment order effects whenever a baseline precedes a treatment.

Christensen (1991) discussed several alternatives for controlling the immediate previous nutritional states of test subjects. For a diet that is to be consumed the next day, for example, the experimenter can request abstinence from eating following the evening meal or can administer a standard meal before the test. Both conditions might require that subjects receive instructions about what else they can or cannot eat or drink. The subjects' "normal" eating is being changed, which sets up potential changes in, for example, eating habit patterns and concern about what is being done to them.

In some cases, Christensen (1991) recommends an elimination or washout phase for 2 to 3 weeks, such as those used in allergy research. They point out that the elimination phase allows for baseline assessment of behavior and permits the full range of the response to manifest itself after reintroduction of the food. A washout phase can clearly show the benefits of avoiding the food and the detriments or advantages of adding the food to the diet. However, this method or approach is likely to influence subjects' expectations. Washout phases must be done with the subjects' permission and cooperation. The subjects must think, for example, that they are doing something to help themselves, something aversive, or something difficult. A washout condition is not like a neutral baseline where one measures "normal" behavior before the introduction of the treatments. The washout condition is a treatment itself that can produce, for instance, withdrawal effects (for caffeine, see Griffiths and Woodson [1988]), varying widely in incidence, severity, and duration. Therefore, subjects must be warned of potential adverse effects, perhaps exacerbates their expectancies. Christensen and colleagues (1991) further point out that the treatment does not always result in prewashout levels.

In addition to attending to dietary baselines, one must also consider the subjects' overall baseline condition. Their prior psychological state could be significant if mood changes are a possible or likely outcome of the dietary treatment. Psychological state could also affect performance on some tasks (e.g., demanding cognitive tasks).

Researchers need to decide whether to consider baseline behavior and mood under conditions of nutrition similar to those that occur in field or nonfield situations. It has been well documented that soldiers in the field, on average, do not eat all of their food. Many eat so little that they lose weight. This issue of underconsumption is dealt with below; the issue here is to highlight the problem of specifying the nutrient status under which the baseline is determined.

Thus, before one even begins a test with a food stimulus, there is a context problem. In what way does one determine baseline performance, and in what way does one examine behavior or mood without any treatment? Many ways of establishing baselines might be as potent as treatments in changing food habits.

### EXPECTANCY

Studies in both humans and animals have repeatedly shown that subjects' expectancies and attributions regarding an "intervention" can dramatically change cognitive, emotional, and behavioral responses. Placebo effects are of critical importance in drug-behavior research. The risk of such attitudinal effects has led investigators to use, for example, double-blind crossover

designs with or that they find a

Although stu- cies, for exam- potential perfor- purpose of this- tions will be p- soldiers face. O- point of view. Rohsenow (198- pharmacologica- the organism eit- of ingestion ar- whether or not

Caffeine is a maintain alertne- Caffeine has b- function as a , double-blind tr- caffeine users ( coffee per day). studies of caffei- have found cle- blood pressure, and eating like (Lick and Boot

Given the p- needed not ju- substances but- situations. For- expectation of- is consumed. A- a military pack- typical underco- attributions wo- agents but also- nutrient suppl- soldiers to obta

designs with or without washout phases to be able to conclude that any effects that they find are due to the drug or nutrient under study.

Although studies designed to control for the cognitive effects of expectancies, for example, are an important and necessary step in the evaluation of potential performance-enhancing substances for the practitioner or, for the purpose of this volume, the U.S. military, the issue is whether these interventions will be potent in both the expected and unexpected real-life scenarios soldiers face. One cannot approach these questions from a purely physiological point of view. For instance, Siegel (1988) in rat models and Marlatt and Rohsenow (1980) in a study of humans have shown that in situations in which pharmacologically active doses of potent, addictive drugs are administered with the organism either expecting or not expecting the substance, the consequences of ingestion are determined more by the organism's expectancy than by whether or not the substance is consumed.

Caffeine is a good example because it is commonly used in an effort to maintain alertness and performance by both civilian and military personnel. Caffeine has been shown to be effective at maintaining alertness and to function as a potent reinforcer (Griffiths and Woodson, 1988). A recent double-blind trial found that withdrawal symptoms can occur in chronic caffeine users even when that use is modest (i.e., about 0.59 l [2.5 cups] of coffee per day). These findings indicate the potency of caffeine. Nonetheless, studies of caffeine (Fillmore and Vogel-Sprott, 1992; Kirsch and Weixel, 1988) have found clear effects of expectancies on measures as diverse as mood, blood pressure, and motor performance. Parallel studies on placebos, alcohol, and eating likewise indicate that expectancies improve behavioral outcomes (Lick and Bootzin, 1975; Marlatt and Rohsenow, 1980; Polivy, 1976).

Given the potential importance of these cognitive mediators, research is needed not just on controlling them to determine the utility of various substances but also on understanding and applying these influences in relevant situations. For example, Cardello and Sawyer (1992) have shown that one's expectation of a food determines its acceptability and is related to how much is consumed. A number of studies have shown that military food and food in a military package are perceived unfavorably, which may explain, in part, the typical underconsumption of rations by soldiers. Studies of expectancies and attributions would help not only in determining the ultimate usefulness of agents but also in developing guidelines for administering the substances (e.g., nutrient supplements within food items versus in pill form) and educating soldiers to obtain the best match of cognitive and physiological forces.

## CONTROLLING THE FOOD, STIMULUS, AND TREATMENT

A number of reviews have highlighted the need for carefully controlled treatment conditions. Kruesi and Rapaport (1986) call for double-blind treatment conditions in which neither the subject nor the experimenter can identify treatment conditions. Blind conditions are essential when the subject or experimenter is likely to have beliefs or expectations about the treatment in particular or the behavioral effects of foods in general. Christensen (1991) even suggests that expectancy can produce behavior that mimics treatment effects under the placebo condition and that one must use continuation of an effect to determine a real treatment effect. Many people claim sensitivity to monosodium glutamate, but very few people actually show an allergic reaction under blind conditions (Meiselman, 1987).

To utilize blind conditions, there must be a treatment substitute (placebo) that is indistinguishable from the treatment. Substituting decaffeinated coffee for caffeinated coffee, for example, appears possible in the short term; but making large dietary changes, like substituting aspartame for sugar in all foods and over an extended period of time, is often not possible (Christensen 1991).

Neither the treatment nor the placebo condition should require the subject to expend more effort. Studies have shown that increased effort affects food choices and intake in both laboratory and natural eating environments (Engell and Hirsch, 1991; Meiselman, et al., 1994). When the effort to obtain a food is increased, selection and, as a result, intake of that food decrease. Moreover, the depressed intake does not necessarily appear to recover quickly or fully when the need for increased effort to obtain a food is removed.

Controlling the treatment often requires placement of the treatment at a certain frequency within the 24-h cycle. The experimenter must decide how often and at what times to deliver the treatment. These times and cycles might not coincide with the subject's normal daily patterns of eating or activity. The issue is whether to allow the subject to control the timing of treatment, or permit the experimenter to test all subjects at the same treatment time.

## PEOPLE

### Individual Differences

It is a truism to say that the effects of a given nutrient or other form of intervention vary from person to person. In broad terms, individual differences in response can be either continuum based (i.e., the dose-response relationship varies across individuals) or category based (i.e., a subset of individuals does not respond to the intervention or some individuals might respond in a paradoxical manner). One needs to determine not only the average response

but also whether the method they may need to intervene is it the cause and the person-specific

Individuals may function differently. However, the real risk with a well-known nutrient is (1979). The mess or is the chance of consuming

Also, nutrients in intake over depending (1979; Nelson). Responses evaluation: variation. (1992a,b) variability. Subjects concluded more than for situations were this focus stable behavior n

but also who responds and to what degree, and one must attempt to determine whether the differences between people are due to their intrinsic characteristics or to other outside influences. In part, this speaks to determining the dosage method that best captures the population of interest. Another assumption that may need to be addressed is person-situation consistency. In other words, if one can reliably describe how a given set of people will respond to the intervention in a particular situation, can one generalize across situations; or is it the case that exposure to physical or psychological stress will change who does and who does not respond? The ultimate goal is to derive a situation- and person-specific dosing system.

Individual differences also can play a mediating role. For example, caffeine may function fairly successfully as an alertness enhancer in most people, or tyrosine might aid in the reduction of stress-based decrements in performance. However, as noted previously, if the relevant source items are not consumed—a real risk with military rations—then the effectiveness or lack thereof is moot. A well-known fact in dietetic studies is that intraindividual variation for most nutrients is as large or larger than interindividual variation (Beaton et al., 1979). Thus, if the agent of interest has a relatively narrow range of effectiveness or is otherwise inflexible regarding its daily required dose, a substantial chance of ineffectiveness arises solely because of soldiers' inconsistencies in consuming source rations or items.

Also of relevance here is that to determine the true daily intake of nutrients in the face of large intraindividual variations, repeated assessment of intake over a number of representative days is required (about 7–14 days depending on the nutrient of interest) (Basiotis et al., 1987; Beaton et al., 1979; Nelson et al., 1989). Similarly, inter- and intravariations in subjects' responses to nutrients could be determined and used to guide the length of evaluations as well as to help investigators understand better the individual variation. Beaton and colleagues (Beaton et al., 1979; Tarasuk and Beaton, 1992a,b) discuss at length the nature and implications of within-subject variability in nutrient intake. A study by Caputo and Mattes (1992) is typical. Subjects were overfed or underfed with carbohydrate or fat. The authors concluded that, although as a group, subjects compensated better for dilutions than for supplementations of energy intake, individual patterns of compensations were highly variable. Likewise, Epstein (1980, 1983) has championed this focus on extended assessment over time as the best approach to obtaining stable behavioral measures. In general, studies on the role of nutrition on behavior might have to be longer than is currently the norm.

### Clinical Disorders

One aspect of individual differences may be represented by individuals with clinical deficiencies or excesses. For example, food and mood (Herman et al., 1987; Leathwood and Pollet, 1982-1983; Morley et al., 1986) are frequently linked, as are mood and performance. Both anxiety and depression have been linked to changes in behavioral activation and food intake. Although the "typical" response to such negative moods—especially at high levels—is a decrement in food intake as well as performance, there are subgroups that tend to show enhanced eating in the face of such stressors. Although clinical disorders per se may not be of central importance to examining the nutrition-performance relationship in the military, such studies do provide a basis for examining the potential role of underlying physiological or psychological states on the relationship.

For example, although both fear and hunger alone can potentiate behavior, when both are present the combination of an aversive and appetitive stimulus can lead to different experimental conclusions, depending on the assessment methodology. As reported by Gray (1987), fear reduces the probability of eating in a hungry subject, but if eating does occur it is more vigorous than that in a nonfearful subject. It is not unreasonable to expect that different mood states will influence the probability of behavioral choices and, in turn, also influence the strength of that behavior once it is chosen.

A more apt example of pathology relates to the impact of jet lag or shift work on performance. This topic has received considerable attention from U.S. Army Research Institute of Environmental Medicine (USARIEM) as well as many other military and civilian research centers (Moline et al., 1990; Tepas, 1990). Circadian rhythms appear to be intimately linked to metabolism, mood, and behavior, and as has been shown repeatedly, disruptions of these rhythms are also disruptive to performance (Armstrong, 1980; Boulos and Terman, 1980; Halberg, 1989). Given the nature of military operations, challenges to circadian rhythms are to be expected, and in instances of rapid deployment, they pose a potentially serious risk. Circadian rhythms may also have a mediating role on food and performance relationships (Mistlberger, 1990). Metabolic research suggests that physiological responses to nutrients vary depending on the time of day when the nutrients are ingested. In addition, the timing of a meal appears to influence some, although not all, biological rhythms. The appropriate type of dietary component and the appropriate dose and schedule of ingestion depend, among other factors, on the schedule in force prior to the disruption, the specifics of the actual disruption, and the feeding and sleep schedule that is possible following the disruption.

Although social sources have learned about how social influences are difficult to manifest in several ways over time of occurrence (e.g., versus influences that rely on verbal modeling and direct or indirect impact to enhance or detract from performance, 1991). Furthermore, social sources or groups. The majority of studies of how much a unit eats a large or small meal. These issues such as unit morale and variables affect unit performance.

A growing body of work (1992) indicates that combat people with whom they are seen with other behavioral colleagues appear to be increases as well, while the data to date do not indicate a determinant of this social behavior.

Studies have also found that sergeant and the soldier have direct effects on consumption. Devens found that soldiers ate more acceptable foods as more acceptable comment and ate most comment and ate appropriate. Similarly, following a 2-week deployment, typically ate their meals or colleagues) consumed who reported that they undifferentiated group (unpublished data, 1991).

Studies with military influences such as mood and behaviors toward food (1991). An interesting aspect of



### Social Influence

Although social sources of influence are pervasive, much remains to be learned about how social factors influence performance and behavior. Social influences are difficult to assess, in part because such influences can be manifested in several ways. Social influences may be categorized in terms of time of occurrence (e.g., influences that occur at the time of measurement versus influences that reflect past experience) and by pathway (e.g., behavioral or verbal modeling and social context) and whether the influences have a direct or indirect impact on what is being measured. Social influences can enhance or detract from concurrent physiological factors (Goldman et al., 1991). Furthermore, social influences can be evaluated in terms of individuals or groups. The majority of food intake research focuses on individuals, as in studies of how much a subject consumes when a confederate eats a relatively large or small meal. The military, however, has a long-standing interest in issues such as unit morale and cohesion and how these group-oriented variables affect unit performance.

A growing body of work (de Castro and Brewer, 1991; Redd and de Castro, 1992) indicates that consumption increases for individuals as the number of people with whom they are eating increases. This social facilitation has been seen with other behaviors (Latane, 1981). The findings of de Castro and colleagues appear to be at least partially explained by the fact that meal length increases as well, while the rate of eating remains nearly the same. However, the data to date do not clearly indicate that increased meal length is the sole determinant of this social facilitation of eating.

Studies have also found that the behaviors or actions of others (e.g., a sergeant and the soldiers under his or her command) can have substantial direct effects on consumption and acceptability. For example, research at Fort Devens found that soldiers ate significantly more rations at a meal and rated foods as more acceptable when their sergeant made a moderate positive comment and ate most of their meal than when he made a moderate negative comment and ate approximately two-thirds of his meal (Engell et al., 1990). Similarly, following a 2-week ration field test, soldiers who reported that they typically ate their meals in a social setting (i.e., with a small group of friends or colleagues) consumed approximately 5 percent more calories than soldiers who reported that they typically ate either alone or in the midst of a large, undifferentiated group of fellow soldiers (F. M. Kramer, U.S. Army Natick, unpublished data, 1991).

Studies with military and civilian personnel have also shown that social influences such as modeling and persuasion can influence attitudes and behaviors toward food (Edelman et al., 1986; Polivy et al., 1979; Smith, 1961). An interesting aspect of that research was that changes in behavior (e.g.,

consumption) and attitudes (e.g., hedonic ratings) were not always consistent with one another (Meiselman et al., 1988). For example, Smith (1961) found that after an experienced U.S. Army sergeant explained to his men the importance of eating unusual foods such as grasshoppers in order to survive and concluded his talk by actually eating a grasshopper, 90 percent of subjects ate one or more grasshoppers; in contrast, hedonic ratings gave no indication of an attitude change. On the other hand, following a cognitive dissonance approach, after soldiers heard a talk by an aloof professional who offered 50 cents to each person who would eat a grasshopper, only 50 percent of subjects actually chose to do so, although hedonic ratings indicated a significant, positive change in attitude. As in other types of research, different measurement methods or different aspects of a phenomenon may yield varying results and conclusions. To be sure of their findings, investigators need to be cognizant of a multitrait-multimethod approach (Campbell and Fiske, 1959).

### CHEMICALS, FOODS, AND MEALS

In a number of human eating research areas, the relevance of stimuli has been discussed (Meiselman, 1992; Spitzer and Rodin, 1981). Studies of nutrition and behavior probably run the range from pure chemical stimuli to entire complex, long-term diets. Different levels of treatment complexity are listed in Table 8-1. Each level of complexity brings with it a set of advantages and a set of disadvantages. Moreover, within each level there can be further distinctions; pure chemical substances can be given in subphysiological doses, physiological doses, or superphysiological doses. Complex meals can be three or more components all on one plate, or they can be multiple courses. Complex diets can involve three meals per day, or they can involve ad libitum consumption of a wide range of foods.

Within the wide range of possible stimuli in food behavior studies, many variables influence food intake habits. For example, Engell (1992) has recently shown that in a meal setting, increased beverage variety leads to increased beverage consumption and increased food consumption. How one constructs laboratory meals and diets in many ways determines the outcomes of studies.

Ultimately, researchers need to know what to serve as an entire diet since that is what all people eat. Although pure substances and simple foods and meals might need to be researched to get the building blocks for diets, can one safely assume that the effects seen in isolation will remain when added to the overall diet? Or, conversely, will dietary components, produce behavioral or mood effects only when added to other dietary components, as Kruesi and Rapaport (1986) suggest?

TABLE 8-1 Level

## Treatment Category

Pure substances

Simple foods

Complex foods

Simple meals

Complex meals

Simple diets

Complex diets

All investigators  
 lence ahead. The  
 ration components  
 rations "would be  
 would be sufficien  
 packaged to withs:  
 tance." This requi  
 than was obtained  
 (Hirsch and Krame  
 present, the long-  
 demands for both

TABLE 8-1 Levels of Treatment Complexity

Treatment Category	Examples
Pure substances	Subphysiological dosage Physiological dosage (Griffiths and Woodson, 1988) Greater than physiological dosage--better enhancement of behavioral effect (Spring et al., 1986)
Simple foods	Pure protein (turkey salad) or pure carbohydrate (bread-like wheat starch) (Spring et al., 1986) Breakfast drink or no food for breakfast (Benton and Sargent, 1992)
Complex foods	Ham casserole as a meal (van Amelsvoort and Weststrate, 1992)
Simple meals	Different breads with cheese, margarine, and coffee or tea (Holm and Bjorck, 1992)
Complex meals	Buffet lunch on 3 alternate days with preload and taste trays (Rolls et al., 1992)
Simple diets	Liquid diet for space travel, 19 weeks, prisoners (Winitz et al., 1965) Two basic meals high and low in energy for 5 days (Caputo and Mattes, 1992)
Complex diets	21-day diet (oat bran, wheat bran intake) in metabolic ward (Anderson et al., 1991) Vending machine diet of entrees, desserts, snacks, and beverages for 7 days (Rising et al., 1992)

All investigators should be clear about the current situation and the challenge ahead. The written military requirement for performance-enhancing ration components (dated July 30, 1992) states that the performance enhancing rations "would be sufficiently stable to withstand long-term storage; they would be sufficiently appealing to ensure consumption; and they would be packaged to withstand standard ration abuses and to promote ration acceptance." This requirement expects more from a performance-enhancing ration than was obtained from the current ration, the Meal Ready to Eat (MRE) (Hirsch and Kramer, 1993; Lester et al., in press; Meiselman et al., 1988). At present, the long-term storage requirements for rations lead to difficult demands for both the developer and the consumer, but these are outside the

topic of this chapter. However, although the Army provides the soldier 3,900 kcal/day for moderate activity in a temperate climate, at present a soldier eats well under 3,000 kcal/day on average and occasionally eats under 2,000 kcal/day, on average.

Many data have been collected from both short-term (Table 8-2) and long-term (Table 8-3) studies of acceptance and consumption of the MREs. Although earlier studies suggested a gradual increase in MRE consumption, this trend has not continued. It is clear that MRE consumption is far more complicated than simply how acceptable the foods are rated.

Since the goal is to enhance performance in combat, one must also question whether a soldier in actual combat would eat more or less than what field training exercise studies have shown. Questionnaire data collected from U.S. Marines who had been in combat indicated that they ate as little as 58 per-

**TABLE 8-2** Prototype Testing (Short-Term Study) of Meal Ready to Eat Rations

Reference	Specific Ration*	Test Duration (days) <sup>†</sup>	Intake (kcal/day)	Acceptance	
				Overall	Main Dish
Askew et al. (1987)	MRE IV	12 (3 MREs/day)	2,282		NA <sup>‡</sup>
Popper et al. (1987)	MRE IV	11 (3 MREs/day)	2,517		5.7
	MRE VII		2,517		6.8
	IMP MRE		2,842	7.86	7.6
Engell et al. (1987)	MRE V	10 (4 MREs/day)	2,733		6.2
Morgan et al. (1988)	MRE VIII	11 (4 MREs/day)	3,217		
Lester et al. (1990)	MRE VIII	3 (4 MREs/day)	2,550		7.03
			2,289		7.39
			2,206		6.56
Edwards et al. (1989)	MRE VI	10 (4 MREs/day)	2,009		6.34
	MRE VIII		2,802		7.33
Lester et al. (1993)	MRE VIII	7 (3 MREs/day)	1,956	5.6	6.8

\* Starting in 1980 the Roman numeral indicates the year of production; for example, MRE IV was produced in 1984.

<sup>†</sup> NA= not available.

<sup>‡</sup> IMP MRE was improved MRE.

**TABLE 8-3**

Reference

Hirsh et al. (1985)

Hirsch and Kramer (1993)<sup>†</sup>

Askew et al.

Thomas et al.<sup>§</sup>

\* Starting in 198 example, MRE I

<sup>†</sup> Student volunte

<sup>‡</sup> Final questionn

<sup>§</sup> Technical Repo

cent of their nor eating more than the same as usual underconsumption worse than the u

At this point, both the food (e.g. situation (e.g., ef time one can act might be able to soldiers, but there

PE

What is the r foods? Researche habits. The milit academia can do interest and inves militarily relevant yet mature. At p measure or how t

TABLE 8-3 Extended Testing of Meal Ready to Eat Rations

Reference	Specific Ration*	Test Duration (days)	Intake (kcal/day)	Acceptance	
				Overall	Main Dish
Hirsh et al. (1985)	MRE IV	34	2,189	7.05	7.05
Hirsch and Kramer (1993) <sup>†</sup>	MRE IV	45	3,149	6.05	
Askew et al.	MRE VI	30	2,782	6.53 <sup>‡</sup>	6.99
Thomas et al. <sup>§</sup>	MRE XII	30	2,441		

\* Starting in 1980 the Roman numeral indicates the year of production; for example, MRE IV was produced in 1984.

<sup>†</sup> Student volunteers.

<sup>‡</sup> Final questionnaire.

<sup>§</sup> Technical Report in preparation.

cent of their normal amount (Popper et al., 1989). Only 3 percent reported eating more than usual on the first combat day, and 29 percent reported eating the same as usual on the first combat day. Therefore, it is highly likely that underconsumption of rations in a real combat situation would be the same or worse than the underconsumption observed in training exercises.

At this point, the ability to predict the food intake of soldiers depends on both the food (e.g., its packaging, palatability, and form [e.g., liquid]) and the situation (e.g., effort required, timing of the meal, and social factors). By the time one can actually use a performance-enhancing ration in the field, one might be able to control and predict its acceptance and consumption by soldiers, but there is much work in this area that has yet to be done.

### PERFORMANCE AND MOOD OUTCOMES

What is the military looking for with regard to performance-enhancing foods? Researchers are interested in what controls human food consumption habits. The military, however, does not have a special need to study what academia can do just as well. Performance-enhancing foods are a subject of interest and investment for the military because it is believed they can enhance militarily relevant behavior. However, the science of soldier performance is not yet mature. At present it is safe to say that researchers do not know what to measure or how to measure it. The measures of physical performance have not

changed much over the years, but there is no way to integrate some of those into an overall physical effectiveness score. Measures of cognitive performance have changed, although integrated measurement of cognitive effectiveness is still far off. Combining physical and cognitive scores has probably never been attempted. Variables such as personality and mood must be factored in as well.

A broader issue in performance enhancement measurement is also one of context. Military thinkers involved with soldiers are necessarily thinking in terms of what capabilities soldiers bring to the battlefield. Recently, a newly proposed Army thrust called the Warrior's Edge focuses on enhancement of a soldier's performance in five capabilities: lethality, mobility, command and control, survivability, and sustainment.

Researchers need to begin to think of how to measure these militarily relevant factors and to demonstrate performance enhancement (or decrements) with these measures. A number of measures of lethality pertaining to firing a rifle exist. Similarly, measures of mobility on foot, either marching or on a treadmill, also exist. Command and control deals with the largely cognitive area of communication and information transfer. The greatest problems are survivability and sustainment. For neither of these is there an established measure.

Perhaps most importantly, researchers need to find ways to combine these measures (and perhaps others) to yield measures of overall soldier effectiveness. The USARIEM has begun some modeling work in this area. The ultimate context for performance-enhancing foods for the military will be the field, and measurement of enhanced soldier performance in the field must be the goal.

Assessment of enhanced soldier performance in the field requires that the enhancements provided by nutrient engineering be of sufficient magnitude to be observable and to make a real difference on the battlefield. The literature in this field is filled with caveats about the sizes of effects. Christensen (1991) distinguish chronic nutritional effects that can produce behavioral changes from acute diet manipulations that produce subtle, nonclinical behavioral changes. He suggests that acute manipulations must be done chronically to observe less subtle behavioral effects. They further warn that the psychological context interacts with chronic effects to produce behavioral changes. This means that short-term tests could produce incorrect estimates of behavioral changes—and might even miss them—and that the psychological context might be a necessary component of such demonstrations.

## CONCLUSIONS

Consideration of context when planning or interpreting the effects of foods on the behaviors or moods of soldiers will require the following changes:

## THE ROLE OF CONTEXT

- Questions will need to plan and conduct the variation of foods and influences), environment, mood and performance

- In order to consider in the following direct foods in diets affect people (e.g., young male, 6- or 12-month period x environment. Then variables into specific

- Investigators will understanding that the

- Anderson, J.W., N.H. Gilins' P.R. Oeltgen  
1991 Lipid responses of Nutr. 54:678-683.
- Armstrong, S.  
1980 A chronometric ap 4:27-53.
- Askew, E.W., I.Munro, M.A. Martin, K. Reynold  
1987 Nutritional status z consuming the Rati a 30-day field train. Institute of Environ
- Askew, E.W., J.R. Claybaug  
1987 Nutrient intakes an ft. altitude consum Mass.: U.S. Army
- Askew, E.W., J.R. Claybaug  
1991 Nutrient intakes an ft. altitude consum Mass.: U.S. Army
- Basiotis, P.P., S.O. Welsh, .  
1987 Number of days of intakes with define
- Beaton, G.H., J. Milner, P. Hewitt, P.V. Grambich, N.  
1979 Sources of variance and interpretation.
- Benton, D., and J. Sargent  
1992 Breakfast, blood gl

ne of those  
performance  
tiveness is  
never been  
l in as well.  
also one of  
thinking in  
ly, a newly  
ncement of  
mmand and

e militarily  
decrements)  
g to firing a  
ing or on a  
ly cognitive  
roblems are  
established

mbine these  
er effective-  
The ultimate  
he field, and  
be the goal.  
ires that the  
magnitude to  
he literature  
nsen (1991)  
hanges from  
ral changes.  
observe less  
gical context  
s means that  
hanges—and  
e a necessary

ffects of foods  
g changes:

• Questions will need to be phrased contextually, and investigators will need to plan and conduct their research contextually. This might involve consideration of foods and meals, people issues (individual differences, social influences), environmental issues (both physical and social) and measures of mood and performance.

• In order to consider context, investigators will need to shift their questions in the following direction: Does carbohydrate (when present in x form in y foods in z diets) affect the behavior (e.g., mobility) or mood (e.g., stress) of people (e.g., young males or soldiers of particular selection criteria) over a 3-, 6-, or 12-month period, as compared with a control condition when tested in x environment. Then investigators will need to translate those contextual variables into specific test conditions.

• Investigators will need to limit their generalizations of results with the understanding that the phenomena might not extend to other contexts.

## REFERENCES

- Anderson, J.W., N.H. Gilinsky, D.A. Deakins, S.F. Smith, D.S. O'Neal, D.W. Dillon, and P.R. Oeltgen  
1991 Lipid responses of hypercholesterolemic men to oat bran and wheat bran. *Am. J. Clin. Nutr.* 54:678-683.
- Armstrong, S.  
1980 A chronometric approach to the study of feeding behavior. *Neurosci. Biobehav. Rev.* 4:27-53.
- Askew, E.W., I.Munro, M.A. Sharp, S. Siegel, R. Popper, M.S. Rose, R.W. Hoyt, J.W. Martin, K. Reynolds, H.R. Lieberman, D.B. Engell, and C.P. Shaw  
1987 Nutritional status and physical and mental performance of special operations soldiers consuming the Ration, Lightweight or the Meal, Ready-to-Eat military field ration during a 30-day field training exercise. Technical Report T7-87. Natick, Mass.: Army Research Institute of Environmental Medicine.
- Askew, E.W., J.R. Claybaugh, S.A. Cucinell, A.J. Young, and E.G. Szeto  
1987 Nutrient intakes and work performance of soldiers during seven days of exercise at 7,200 ft. altitude consuming the meal, Ready-to-Eat ration. Technical Report T3-87. Natick, Mass.: U.S. Army Research Institute of Environmental Medicine.
- Askew, E.W., J.R. Claybaugh, S.A. Cucinell, A.J. Young, and E.G. Szeto  
1991 Nutrient intakes and work performance of soldiers during seven days of exercise at 7,200 ft. altitude consuming the Meal, Ready-to-Eat ration. Technical Report T3-87. Natick, Mass.: U.S. Army Research Institute of Environmental Medicine.
- Basiotis, P.P., S.O. Welsh, F.J. Cronin, J.L. Kelsay, and W. Mertz  
1987 Number of days of food intake records required to estimate individual and group nutrient intakes with defined confidence. *J. Nutr.* 117:1638-1641.
- Beaton, G.H., J. Milner, P. Corey, V.M. McGuire, M. Cousins, E. Stewart, M. de Ramos, D. Hewitt, P.V. Grambich, N. Kassim, and J.A. Little  
1979 Sources of variance in 24-hour dietary recall data: Implications for nutrition study design and interpretation. *Am. J. Clin. Nutr.* 32:2546-2559.
- Benton, D., and J. Sargent  
1992 Breakfast, blood glucose and memory. *Biol. Psych.* 33:207-210.

- Birch, L.L., J. Billman, and S.S. Richards  
1984 Time of day influences food acceptability. *Appetite* 5:109-116.
- Blundell, J.E., and A.J. Hill  
1986 Paradoxical aspects of an intense sweetener (aspartame) on appetite. *Lancet* 1:1092-1093 (letter).
- Blundell, J.E., P.J. Rogers, and A.J. Hill  
1988 Uncoupling sweetness and calories: Methodological aspects of laboratory studies on appetite control. *Appetite* 11(Suppl.):54-61.
- Boulos, Z., and M. Terman  
1980 Food availability and daily biological rhythms. *Neurosci. Biobehav. Rev.* 4:119-131.
- Campbell, D.T., and D. Fiske  
1959 Convergent and discriminant validation by the multitrait-multimethod matrix. *Psychological Bull.* 56:81-105.
- Campbell, D.T., and J.C. Stanley  
1966 *Experimental and Quasi-Experimental Designs for Research*. Skokie, Ill.: Rand McNally.
- Caputo, F.A., and R.D. Mattes  
1992 Human dietary responses to covert manipulations of energy, fat, and carbohydrate in a midday meal. *Am. J. Clin. Nutr.* 56:36-43.
- Cardello, A.V., and F.M. Sawyer  
1992 Effects of disconfirmed consumer expectations on food acceptability. *J. Sensory Studies* 7:253-277.
- Christensen, L., J. Miller, and D. Johnson  
1991 Issues in the design of studies investigating the behavioral concomitants of foods. *J. Consult. and Clin. Psych.* 6:874-882.
- Collier, G.  
1989 The Economics of Hunger. Thirst, Satiety and Regulation. *Ann. N.Y. Acad. Science.*, 575:136-154.
- de Castro, J.M., and E.M. Brewer  
1991 The amount eaten in meals by humans is a power function of the number of people present. *Physiol. Behav.* 51:121-125.
- Edelman, B., D. Engell, P. Bronstein, and E. Hirsch  
1986 Environmental effects on the intake of overweight and normal-weight men. *Appetite* 7:71-83.
- Edwards, J.S.A., D.E. Roberts, T.E. Morgan, and L.S. Lester  
1989 An evaluation of the nutritional intake and acceptability of the Meal, Ready-to-Eat consumed with and without a Supplemental pack in a cold environment. Technical Report T18-89. Natick, Mass.: U.S. Army Research Institute of Environmental Medicine.
- Engell, D.  
1992 Sensory influences on meal consumption. Rose Marie Pangborn Memorial Symposium. August 2-6, Jarvenpaa, Finland.
- Engell, D.B., D.E. Roberts, E.W. Askew, M.S. Rose, J. Buchbinder, and M.A. Sharpe  
1987 Evaluation of the Ration Cold Weather during a 10-day cold weather field training exercise. NATICK Technical Report TR-87/030. Natick, Mass.: U.S. Army Natick Research, Development and Engineering Center.
- Engell, D., F.M. Kramer, S. Luther, and S. Adams  
1990 The effect of social influences on food intake. Paper presented at the Society for Nutrition Education 1990 Annual Meeting, Anaheim, Calif.
- Epstein, S.  
1980 The stability of behavior. II. Implications for psychological research. *Am. Psychologist* 35:790-806.

- 1983 Aggregation and 51:360-392.
- Fillmore, M., and M. Vo  
1992 Expected effect placebo. *Psych*
- Goldman, S.J., C.P. Herm  
1991 Is the effect of by hunger? *A*
- Gray, J.A.  
1987 The Psychology  
Griffiths, R.R., and P.P.  
1988 Reinforcing effe  
Halberg, F.  
1989 Some aspects of *J. Nutr.* 119:333
- Herman, C.P., J. Polivy,  
1987 Anxiety, hunger  
Hirsch, E., and F.M. Kra  
1993 Situational influe  
of the Committee  
Board, Institute
- Hirsch, E., H. L. Meiselm  
M. Fox, S. McN  
1985 The effects of  
Technical Repo  
Development an
- Holm, J., and I. Bjorck  
1992 Bioavailability o  
responses in heal  
*Nutr.* 55:420-42
- Kirsch, I., and L.J. Weixe  
1988 Double-blind v  
102:319-323.
- Kramer, F.M., K. Rock, :  
1992 Effects of time o  
and mid-day. *Ap*
- Kruesi, M.J., and J.L. Raj  
1986 Diet and human  
Review of Nutrit  
Annual Reviews
- Latane, B.  
1981 The psychology  
Leathwood, P.D., and P.  
1982- Diet-induced mo  
1983 17:147-154.
- Lester, L.S., F.M. Kramer  
1990 Evaluation of the  
consumption of t  
TR-90/008L. Nat  
Center.



- 1983 Aggregation and beyond: Some basic issues on the prediction of behavior. *J. Personality* 51:360-392.
- Fillmore, M., and M. Vogel-Sprott  
1992 Expected effect of caffeine on motor performance predicts the type of response to placebo. *Psychopharmacology* 106:209-214.
- Goldman, S.J., C.P. Herman, and J. Polivy  
1991 Is the effect of a social model on eating attenuated by hunger? *Appetite* 17:129-140.
- Gray, J.A.  
1987 *The Psychology of Fear and Stress*. Cambridge: Cambridge University Press.
- Griffiths, R.R., and P.P. Woodson  
1988 Reinforcing effects of caffeine in humans. *J. Pharmacol. Exp. Ther.* 246:21-29.
- Halberg, F.  
1989 Some aspects of the chronobiology of nutrition: More work is needed on "when to eat." *J. Nutr.* 119:333-343.
- Herman, C.P., J. Polivy, C.N. Lank, and T.F. Heatherton  
1987 Anxiety, hunger, and eating behavior. *J. Abnormal Psychol.* 96:264-269.
- Hirsch, E., and F.M. Kramer  
1993 Situational influences on food intake. In *Nutrition for Work in Hot Environments*. Report of the Committee on Military Nutrition Research, B.M. Marriott, ed., Food and Nutrition Board, Institute of Medicine. Washington, D.C.: National Academy Press.
- Hirsch, E., H. L. Meiselman, R. D. Popper, G. Smits, B. Jezior, I. Lichton, N. Wenkam, J. Burt, M. Fox, S. McNutt, T.N. Thiele, and O. Direge  
1985 The effects of prolonged feeding Meal, Ready-to-Eat (MRE) operational rations. Technical Report Natick TR-85/035. Natick, Mass.: U.S. Army Natick Research, Development and Engineering Center.
- Holm, J., and I. Bjorck  
1992 Bioavailability of starch in various wheat-based bread products: Evaluation of metabolic responses in healthy subjects and rate and extent of in vitro starch digestion. *Am. J. Clin. Nutr.* 55:420-429.
- Kirsch, I., and L.J. Weixel  
1988 Double-blind versus deceptive administration of a placebo. *Behav. Neurosci.* 102:319-323.
- Kramer, F.M., K. Rock, and D. Engell  
1992 Effects of time of day and appropriateness on food intake and hedonic ratings at morning and mid-day. *Appetite* 18:1-13.
- Kruesi, M.J., and J.L. Rapaport  
1986 Diet and human behavior: How much do they affect each other? Pp. 113-130 in *Annual Review of Nutrition*, R. E. Olson, E. Beutler, and H. P. Broquist, eds. Palo Alto, Calif.: Annual Reviews Inc.
- Latane, B.  
1981 The psychology of social impact. *Am. Psychologist* 36:343-355.
- Leathwood, P.D., and P. Pollet  
1982- Diet-induced mood changes in normal populations. *J. Psychiat. Res.*  
1983 17:147-154.
- Lester, L.S., F.M. Kramer, J. Edinberg, S. Mutter, and D.B. Engell  
1990 Evaluation of the Canteen cup stand and ration heater pad: effects on acceptability and consumption of the Meal, Ready-to-Eat in a cold weather environment. Technical Report TR-90/008L. Natick, Mass.: U.S. Army Natick Research, Development and Engineering Center.

- Lester, L.S., L.L. Leshner, M. Salomon, D. B. Engell, S.L. Dewey, J. Ward, C. Thomas, and J. Kalick  
 1993 Nutritional and hedonic consequences of consuming the MRE VIII or the SEP MRE. Technical Report Natick TR-93/015. Natick, Mass.: U.S. Army Natick Research Development and Engineering Center.
- Lick, J., and R. Bootzin  
 1975 Expectancy factors in the treatment of fear: Methodological and theoretical issues. *Psychological Bull.* 82:917-931.
- Marlatt, G.A., and D.J. Rohsenow  
 1980 Cognitive processes in alcohol use: Expectancy and the balanced placebo design. Pp. 159-199 in *Advances in Substance Abuse*, Vol. 1, N. K. Mello, ed. Greenwich, Conn.: JAI Press.
- Meiselman, H.L.  
 1987 Use of sensory evaluation and food habits data in product development. Pp. 307-326 In *Umami: A Basic Taste* Y. Kawamura and M. Kare, eds. New York: Marcel Dekker.  
 1992 Methodology and theory in human eating research. *Appetite* 19:49-55.  
 1994 Identifying and demonstrating consumer acceptance variables in a natural eating environment. In *Proceedings of Food Preservation 2000*, I. Taub and R. Bell, eds. Washington, D.C.: National Academy Press.
- Meiselman, H.L., E.S. Hirsch, and R.D. Popper  
 1988 Sensory, hedonic and situational factors in food acceptance and consumption. Pp. 77-87 in *Food Acceptability*, D. M. H. Thomson, ed. London: Elsevier Applied Science.
- Mistlberger, R.  
 1990 Circadian pitfalls in experimental designs employing food restrictions. *Psychobiology* 18:23-29.
- Moline, M.L., C.P. Pollack, S. Zendell, L.S. Lester, C.A. Salter, and E. Hirsch  
 1990 A Laboratory Study of the Effects of Diet and Bright Light Countermeasures to Jet Lag. Technical Report Natick TR-90/024. Natick, Mass.: U.S. Army Natick Research, Development and Engineering Center.
- Morgan, T.E., L.A. Hodges, D. Schillig, R.W. Hoyt, E.J. Iwanyk, G. McAnich, T.C. Wells, and E.W. Askew  
 1988 A comparison of the Meal, Ready-to-Eat, Ration Cold Weather and Ration, Lightweight nutrient intakes during moderate altitude cold weather field training operations. Technical Report T5-89. Natick, Mass: U.S. Army Research Institute of Environmental Medicine.
- Morley, J.E., A.S. Levine, and M.L. Willenbring  
 1986 Stress-induced feeding disorders. Pp. 71-99 in *Pharmacology of Eating Disorders: Theoretical and Clinical Developments*, M. O. Carruba and J. E. Blundell, eds. New York: Raven Press.
- Nelson, M., A.E. Black, J.A. Morris, and T.J. Cole  
 1989 Between- and within-subject variations in nutrient intake from infancy to old age: Estimating the number of days required to rank dietary intakes with desired precision. *Am. J. Clin. Nutr.* 50:155-167.
- Polivy, J.  
 1976 Perception of calories and regulation of intake in restrained and unrestrained subjects. *Addict. Beh.* 1:237-243.
- Polivy, J., C.P. Herman, J.C. Younger, and B. Erskine  
 1979 Effects of a model on eating behavior: The induction of a restrained eating style. *J. Personality.* 47:100-117.
- Popper, R., E. Hirsch  
 1987 Field Evaluation of the MRE VIII. Technical Report Natick TR-87/027. Natick, Mass.: U.S. Army Natick Research Development and Engineering Center.
- Popper, R., G. Smits  
 1989 Eating in the field. *Appetite* 13:1-10.
- Redd, M., and J.M. J.  
 1992 Social facilitation of eating behavior. *Appetite* 19:749-754.
- Rising, R., S. Alger  
 1992 Food intake and energy expenditure. *Am. J. Clin. Nutr.* 55:1-10.
- Rolls, B.J.  
 1991 Effects of food intake on energy expenditure. *J. Clin. Nutr.* 52:1-10.
- Rolls, B.J., A.E. Anderson  
 1992 Food intake and energy expenditure. *Clin. Nutr.* 11:1-10.
- Rose, M.S. and D.E.  
 1987 Effect of A-100 on food intake. Report T2-8.
- Schutz, H.G.  
 1988 Beyond the energy balance equation. *Am. J. Clin. Nutr.* 48:115-134.
- Siegel, S.  
 1988 State-dependent learning. *Psychol. Rev.* 95:1-24.
- Smith, E.E.  
 1961 The power of suggestion. *Psychol. Rev.* 68:1-10.
- Spitzer, L., and J. R.  
 1981 Human eating behavior. *Psychol. Rev.* 88:1-10.
- Spring, B. J., H.R. Li  
 1986 Effects of cold weather on food intake. *Am. J. Clin. Nutr.* 43:1-10.
- Tarasuk, V., and G.F.  
 1992a Day-to-day variation in food intake. *Am. J. Clin. Nutr.* 55:1-10.  
 1992b Statistical variation in food intake. *Am. J. Clin. Nutr.* 55:1-10.
- Tepas, D.I.  
 1990 Do eating and drinking habits affect energy balance? *Am. J. Clin. Nutr.* 52:203-211.
- van Amelsvoort, J.M.  
 1992 Amylose- and amylose-free diets. *Am. J. Clin. Nutr.* 55:1-10.
- Winitz, M., J. Graff,  
 1965 Evaluation of the MRE VIII. Technical Report Natick TR-65/027. Natick, Mass.: U.S. Army Natick Research Development and Engineering Center.

- Popper, R., E. Hirsch, L. Leshner, D. Engell, B. Jezior, B. Bell, and W.T. Matthew  
1987 Field Evaluation of Improved MRE, MRE VII, and MRE IV. NATICK Technical Report TR-87/027. Natick, Mass.: U.S. Army Natick Research, Development and Engineering Center.
- Popper, R., G. Smits, H.L. Meiselman, and E. Hirsch  
1989 Eating in combat: A survey of U.S. Marines. *Milit. Med.* 154:619-623.
- Redd, M., and J.M. de Castro  
1992 Social facilitation of eating: Effects of social instruction on food intake. *Physiol. Behav.* 52:749-754.
- Rising, R., S. Alger, V. Boyce, H. Seagle, R. Ferraro, A.M. Fontvieille, and E. Ravussin  
1992 Food intake measured by an automated food-selection system: Relationship to energy expenditure. *Am. J. Clin. Nutr.* 55:343-349.
- Rolls, B.J.  
1991 Effects of intense sweeteners on hunger, food intake, and body weight: A review. *Am. J. Clin. Nutr.* 53:872-878.
- Rolls, B.J., A.E. Andersen, T.H. Moran, A.L. McNelis, H.C. Baier, and I.C. Fedoroff  
1992 Food intake, hunger, and satiety after preloads in women with eating disorders. *Am. J. Clin. Nutr.* 55:1093-1103.
- Rose, M.S. and D.E. Carlson  
1987 Effect of A-ration meals on body weight during sustained field operations. Technical Report T2-87. Natick, Mass.: U.S. Army Research Institute of Environmental Medicine.
- Schutz, H.G.  
1988 Beyond preference: Appropriateness as a measure of contextual acceptance of food. Pp. 115-134 in *Food Acceptability*, D.M.H. Thomson, ed. London: Elsevier Applied Science.
- Siegel, S.  
1988 State-dependent learning and morphine tolerance. *Behav. Neurosci.* 102:228-232.
- Smith, E.E.  
1961 The power of dissonance techniques to change attitudes. *Public Opinion Q.* 25:626-639.
- Spitzer, L., and J. Rodin  
1981 Human eating behavior: A critical review of studies in normal weight and overweight individuals. *Appetite* 2:293-329.
- Spring, B.J., H.R. Lieberman, G. Swope, and G.S. Griffiths  
1986 Effects of carbohydrates on mood and behavior. Pp. 51-60 in *Nutrition Reviews*, R. E. Olson, ed. Washington, D.C.: Nutrition Foundation.
- Tarasuk, V., and G.H. Beaton  
1992a Day-to-day variation in energy and nutrient intake: Evidence of individuality in eating behaviors? *Appetite* 18:43-54.  
1992b Statistical estimation of dietary parameters: Implications of patterns in within-subject variation—a case study of sampling strategies. *Am. J. Clin. Nutr.* 55:22-27.
- Tepas, D.I.  
1990 Do eating and drinking schedules interact with work schedule variables? *Work Stress* 4:203-211.
- van Amelsvoort, J.M.M., and J.A. Weststrate  
1992 Amylose-amylopectin ratio in a meal affects postprandial variables in male volunteers. *Am. J. Clin. Nutr.* 55:712-718.
- Winitz, M., J. Graff, N. Gallagher, A. Narkin, and D.A. Seedman  
1965 Evaluation of chemical diets as nutrition for man-in-space. *Nature* 205:742-743.

## DISCUSSION

DAVID D. SCHNAKENBERG: Herb, I want to compliment you on your thought-provoking paper. You have brought out issues that you and I used to talk about.

I would like to add one point to one study that you cited to me, but I feel it was done by Rose and Carlson (1987) on sustained artillery operations. It was serendipity, the study that was done—7 days of continuous artillery operations with very little sleep to see how the performance of these artillerymen would be degraded. The plan was to feed them MREs [Meals Ready to Eat] three times a day. However, the bugs crawled out of some plant in San Antonio, and the Secretary of the Army put a total moratorium throughout the world that "thou shalt not" eat any more MREs until we find the problem in the production plant.

That happened 2 days before the planned study. There was a question: Should they cancel the study or do something else? They could not feed them MREs. So they took hot meals from the dining hall and carried them out to the field, and the cooks prepared and laid out the food for them. They had to stop the action briefly and let the people sit down and eat the meals and then continue on.

That was the study in which soldiers in the field ate 3,700 calories per day over a 7-day period; they lost no water and did not become dehydrated. Much to the frustration I think of Dr. (James) Vogel, they had no performance decrements.

DAVID D. SCHNAKENBERG: So I think that is one example: If there is nothing physiologically or psychologically wrong with a soldier when he goes outside, if you give him something that he likes to eat and in a situation which gives him time to eat it, he will eat to maintain energy balance. However, these factors are often not all in place in the other field studies that have been undertaken.

ROBERT O. NESHEIM: In every one of the MRE tests that we have evaluated as a committee, we have always had that issue that the troops do not eat enough to maintain their weight. The question is, why don't they? I think the context of the situation is a big part of it.

G. RICHARD JANSEN: That gap you mentioned at the end between, say, the real-world situation and experimental protocols, is a real issue.

HERBERT MEISELMAN: I think the focus groups which Natick has conducted might show you some of the things to look at, but you are going to

have to develop some developers, logistici

ROBERT O. NESHEIM: problems that I deal many years ago. W degrees of acceptab products, but somet they would fall flat

It is my underst doing less and less doing more and mor overcome some of t

JOHANNA DWYER: presentations this m revisit this issue of c just now about that from some of the sl very important.

The question I do it? Similarly, doe or are there things tl the next 2 days with

JOEL GRINKER: 1 in England, and the

It certainly is i rhythms because the confuses things. Th

Also, I think Dr the effects of differ

I have another c importance of eating addressed food acce meats, they did expu somebody who pose acceptance than if t

EDWARD HORTON: presentation, at the p and the importance

have to develop something very quantitative after that in order to aid the ration developers, logisticians, and nutritionists..

ROBERT O. NESHEIM: I was reminded, as you were talking, about the problems that I dealt with in a food lab at the Quaker Oats Corporation many, many years ago. We would have very good tests on products that had high degrees of acceptability. They compared very favorably against competitive products, but sometimes you would put them out in a test marketplace and they would fall flat on their face.

It is my understanding that more and more of the food research labs are doing less and less of a detailed product evaluation within the labs, but are doing more and more of it out in the field, simply because they are trying to overcome some of that situation.

JOHANNA DWYER: I just wanted to compliment everybody who made presentations this morning because I thought they were fascinating. I want to revisit this issue of continuity. It seems from David Schnakenberg's statement just now about that experiment where they put the hot foods out, and then from some of the sleep data we heard earlier, this continuity issue seems to be very important.

The question I have is, is it possible to sleep in a tank; is there a way to do it? Similarly, does the context of the combat situation simply preclude this, or are there things that we can consider? What are our constraints as we work the next 2 days with respect to this issue?

JOEL GRINKER: That has been documented by behavioral tests by Dr. Smith in England, and there is no question that there is a decline in intake.

It certainly is important, and it gets complicated in terms of circadian rhythms because the circadian rhythm also has a 12-hour cycle, so that sort of confuses things. That may be part of the cause of that 12-hour cycle.

Also, I think Dr. Bonnie Spring will have some things to say later about the effects of different food on the amount of behavioral change.

I have another comment. I was reminded by the sergeants explaining the importance of eating grasshoppers, about the studies during World War II that addressed food acceptance, in order to get the American public to eat organ meats, they did experiments in which they suggested that if, in fact, they had somebody who posed as an expert—the tests showed—that they got a greater acceptance than if they just told people that this is good for them.

EDWARD HORTON: I am fascinated, as I have listened to this morning's presentation, at the parallelism between sleep deprivation and food deprivation and the importance of the context. If you put a person in a context where he

cannot get sleep for very long or you put him in a context where he cannot sit down and enjoy a good hot meal, you end up with these deprivation syndromes. There seems to be a lot of parallelism.

I was wondering if people have looked at the recovery from these. After you have been through a sleep deprivation, do you sleep like a babe after a while? After you have been through a period of food deprivation, do you go back and chow down and make up your energy losses and so forth? Looking at periodic recovery, are there ways to enhance recovery after the fact?

HERBERT MEISELMAN: I think part of our problem is that the more we study this, the more we realize that our tests should be longer and longer in the field. There is a real problem with keeping troops in the field in a training exercise for 4, 6, or 8 weeks and with the associated costs. I think we have thought of what you are suggesting, but it is very difficult to do; it is very long-term testing.

EDWARD HORTON: There are two questions. One is adaptation; adaptations occur over time. The other is recovery and thinking of something like the exercise physiology test where you basically give little rest periods between tests.

HERBERT MEISELMAN: In our first test in 1982, we were positive that the troops would find the diet very monotonous. We predicted that their food acceptance ratings would go down over time, an example of classic monotony.

In fact, what happened is that the troop ratings of the foods stayed absolutely flat for 35 days. But over time they ate less and less and less, so that by the end of the test they were eating considerably less than at the beginning but were not aware of it.

So our simple model of this—that the diet would be monotonous and they would not eat it—did not fit. The diet, in effect, was monotonous if you view it from an intake measure but not from a rating measure.

II

IN PART IV T  
ment of the ba  
Speakers were  
areas. Each spe  
field of expertis  
to provide copi  
before the work  
were asked to r  
with their reco  
question and an  
are included at  
Reviewed  
administration  
responses to mi  
physiological a  
performance un